

WORLD LEARNING PROGRAM HEALTH GUIDELINES & REQUIREMENTS: CHILE

BRIEF SUMMARY

Prior to Departure:

- Schedule an appointment with your health care provider or a travel medicine clinic at least 8 weeks prior to departure. Please note that some vaccines are only available at travel medicine clinics.
- Take an adequate quantity of all your prescription and non-prescription medications to last your entire stay.
- Ensure you are up to date on all routine vaccines. Routine vaccines that should be up to date prior to traveling include: Covid 19, varicella (chicken pox), hepatitis A & B, HPV, influenza, meningococcal, measles/mumps/rubella (MMR), polio, and tetanus/diphtheria/pertussis (Tdap).
- Discuss your specific travel itinerary (including various in-country locations you plan to visit) with your health care provider. We recommend you ask about and discuss:
 - Are you up to date on your routine vaccines including all boosters?
 - Are there additional vaccines that are recommended? These may include but are not limited to:
 - Typhoid
 - Plan for the prevention and treatment of traveler's diarrhea.
- World Learning recommends that the International Certificate of Vaccination or Prophylaxis (ICVP,) also known as the yellow WHO card, along with a copy of your immunization record should be kept with your passport so that it is easily accessible.

While You Are in Chile:

- Remember motor vehicle accidents remain the biggest risk for international travelers. Ride in the back seat if possible, wear seatbelts, and do not be afraid to ask your driver to slow down.
- Take preventive measures to avoid mosquito and other insect bites.

- Take measures to prevent traveler's diarrhea.
- Do not travel alone and maintain an awareness of your surroundings at all times.
- Use condoms if you are sexually active.
- While traveling abroad, minimize high-risk behaviors.
- Notify your Program staff if you become ill.

IMMUNIZATIONS

Immunizations are recommended to protect your health and well-being by building up your immune defenses against specific prevalent diseases. The decision to receive each vaccine should be made with your primary care provider or Travel Medicine clinic provider.

Plan ahead at least 8 weeks since some immunizations require more than one dose for effectiveness. SIT recommends that you bring a copy of your immunization record with you and insert this document in your passport so that it is easily accessible.

World Learning RECOMMENDS participants are up to date with each of the following vaccinations including initial series and boosters:

- COVID-19
- Hepatitis A
- Hepatitis B
- Meningococcal (meningitis)
- MMR (measles, mumps, rubella)
- Tetanus, diphtheria, pertussis (Tdap)
- Varicella (chicken pox)
- Polio
- Typhoid
 - Recommended for most travelers in Chile, especially those staying with friends or relatives or visiting smaller cities or rural areas.

GENERAL INFORMATION

Maintaining good health and safety is a critical element of your successful participation in any World Learning program. While traveling abroad students should always use caution in unfamiliar places and circumstances. Access to emergency care abroad may not be what you are used to at home. A review of health, safety and security information will be provided to students upon arrival in the host country.

To protect your health in Chile, you may need certain pre-departure immunizations followed by reasonable health precautions while in the country. The following health guidelines and requirements are based on years of experience and the current recommendations from the US Centers for Disease Control and Prevention. They are designed to inform you of health concerns that may be present in Chile.

Although no information sheet can address every conceivable contingency, the following health guidelines and requirements are an attempt to provide you with a standard, which if followed, should optimize good health during your stay abroad.

You may find that local customs and practice, as well as varying US health care providers' approaches, at times conflict with these guidelines. It is essential that you review these health guidelines and requirements with your health care provider and discuss individual issues such as pre-existing medical problems and allergies to specific drugs. Any further questions or concerns should be directed to the US Centers for Disease Control and Prevention (CDC)- www.cdc.gov/travel - or to your own health care provider.

PREVENTION OF INSECT BORNE ILLNESS

Personal Protective Measures (for mosquitos and other insects):

For all the diseases listed below, bug exposure and bite prevention are crucial to reducing the risks to travelers' health. We recommend you take the following steps.

- Prevent bug exposure and bites:
 - Wear long sleeves, long pants, shoes, and hats to minimize exposed skin.
 - Wear clothing and shoes treated with the repellent permethrin. (Permethrin is not for use directly on skin.)
- Choose an appropriate insect repellent and use it regularly:
- Protection against multiple bugs (mosquitos, ticks, flies, etc.): The CDC recommends a repellent which contains at least 20% DEET.
- Protection against mosquitoes only: Repellents other than DEET protect against mosquitos but may not be as effective against other bugs:
 - Picaridin (also known as KBR 3023, Bayrepel and icaridin)
 - Oil of lemon eucalyptus (OLE) or para- menthane-diol (PMD)
 - IR3535 (SkinSoSoft)
 - 2-undecanone (methyl nonyl ketone)
- Always use repellents as directed.
- Please consult your healthcare provider if you have any health concerns regarding bug repellants. (See recommendations above.)
- Avoid transiting tall grass, shrubs, or woody areas and check for ticks afterward.
- Consider using a mosquito net while sleeping.
- Check your entire body for ticks after outdoor activity.
- If you experience symptoms after the program, please consult your healthcare provider and be sure to tell them about your travel.

WORLD LEARNING PROGRAM HEALTH GUIDELINES & REQUIREMENTS: CHILE

For those programs going to Argentina, please note the risk of Dengue and Chaga's disease...

Dengue:

Dengue viruses are spread to people through the bite of an infected mosquito, primarily during the daytime. Dengue symptoms usually start within a few days of being bitten but can take up to 2 weeks to develop. Not everyone who gets infected with dengue will feel sick. Symptoms can be mild or severe and can include fever with nausea, vomiting, rash, headache, eye pain, joint and muscle pain. In severe cases, dengue can cause shock, internal bleeding, and even death. Travelers can protect themselves from dengue by following the personal protective measures to avoid mosquito bites as listed above.

Chaga's Disease:

American trypanosomiasis, also known as Chaga's disease, is caused by a parasite. This parasite lives in the poop of triatomine (kissing) bugs. The bug often poops when it bites. Most people get Chaga's disease by unknowingly rubbing triatomine bug poop into the bug bite when they scratch the bite. People may also accidentally get triatomine bug poop in their eyes and mouth. A few people experience swelling around the bite area. While most people never have symptoms, they can remain infected throughout their lives. About 1 out of 3 people who get Chaga's disease will develop more serious symptoms later in life, including heart conditions or gastrointestinal problems. These later symptoms can occur whether a person has early symptoms or not. Chaga's disease is spread by triatomine bugs in parts of Mexico, Central America, and South America, especially rural areas. Fortunately, most travelers to these areas don't need to worry about Chaga's disease. Travelers who sleep outdoors or stay in poorly constructed housing are more likely to get bitten by triatomine bugs. Travelers can protect themselves by following the personal protective measures listed above.

PREVENTION OF FOOD/WATER BORNE ILLNESS

Diarrhea Producing Infections:

"Traveler's diarrhea" is the most common form of travel related illness. In otherwise healthy adults, diarrhea is rarely serious or life-threatening, but it can make a trip very unpleasant. The diarrhea may last several days and is characterized by watery, non-bloody bowel movements. Traveler's diarrhea usually requires no treatment other than fluid replacement by drinking plenty of water or using an Oral Rehydration Solution (ORS) to prevent dehydration. ORS can be purchased as a prepackaged solution. Antidiarrheals such as Imodium or Lomotil may be used short-term in some circumstances as long as the traveler stays well hydrated and has no bloody stools or abdominal pain. Pepto Bismol can be used to treat diarrhea, but large quantities are required. Antibiotics may be indicated for more severe cases of traveler's diarrhea especially if the traveler has fever or stools with blood or mucous. Ask your health care provider before your program begins for their recommendations regarding traveler's diarrhea.

More protracted and disabling diarrheal illnesses may be due to giardiasis, dysentery, cholera, and typhoid. Other diseases such as hepatitis A are also transmitted via contaminated food and water. These infections as well as "traveler's diarrhea" are often caused by contaminated food and water.

Tap water is generally safe to drink and cook with in Santiago, Chile; Buenos Aires, Argentina and other major cities and it is usually safe to eat fresh fruits, vegetables, and salads. Food and water safety, however, cannot be guaranteed from street food purchases and the water quality is variable in rural areas. If you are unsure of the food and water quality in the area you are visiting, check with a reliable source before using. For program excursions and rural areas, take all the recommended precautions and stick to safe food and water habits including the following:

WORLD LEARNING PROGRAM HEALTH GUIDELINES & REQUIREMENTS: CHILE

DO:

- WASH your hands scrupulously with non-contaminated water and soap before eating.
- DRINK:
 - Bottled water that is sealed
 - Water that has been disinfected
 - Ice made with bottled or disinfected water
 - Carbonated drinks
 - Hot coffee or tea
 - Pasteurized milk
- PURIFY YOUR WATER if it is not bottled or from a reliable source. Options for purification include...
 - Boiling: bring to a full rolling boil for 1 min (3 min at elevations over 6,500 ft)
 - Disinfectants such as iodide and chlorine dioxide (follow the manufacturer's instructions)
 - Filtration devices and UV light (follow the manufacturer's instructions)
- DO EAT:
 - Food that is cooked and served hot
 - Hard-cooked eggs
 - Fruits and vegetables you have washed in clean water or peeled yourself
 - Pasteurized dairy products

DON'T:

- DO NOT DRINK:
 - Tap or well water
 - Ice made with tap or well water
 - Drinks made with tap or well water (such as reconstituted juice)
 - Unpasteurized milk
- DO NOT EAT:

- Food served at room temperature
- Food from street vendors
- Raw or soft-cooked (runny) eggs
- Raw or undercooked (rare) meat or fish
- Unwashed or unpeeled raw fruits and vegetables
- Unpasteurized dairy products
- Bushmeat" (monkeys, bats, or other wild game)

There may be times when refusing an offer of food or beverage, even a drink with ice or avoiding a salad will be considered rude. You must decide for yourself, but polite refusals, thought out in advance, are often handy. Discuss these alternatives with your Program staff.

Hepatitis A:

Hepatitis A is a highly contagious virus that causes liver inflammation. It is most commonly spread through contaminated food and water and is best avoided by practicing safe food and water habits as listed above.

Typhoid:

Typhoid is an infection caused by the salmonella bacterium. People infected with these bacteria can spread them to others. This typically happens when an infected person uses the bathroom and does not wash their hands. The bacteria can stay on their hands and contaminate everything that the person touches, including food and drinks. In countries with poor sanitation, the water used to rinse and prepare food and beverages, including tap water, can also be contaminated with these bacteria. Travelers who eat foods or drink beverages contaminated with these bacteria can then get sick. People with typhoid may have a fever that can be as high as 103 to 104°F (39 to 40°C). They also may have weakness, stomach pain, headache, diarrhea or constipation, cough, and loss of appetite. Antibiotics are used for treatment. To prevent the infection, follow safe food and water habits as listed above and talk with your health care provider about getting the vaccine prior to traveling.

WORLD LEARNING PROGRAM HEALTH GUIDELINES & REQUIREMENTS: CHILE

A Note on Swimming...

Avoid swimming or wading in fresh water. Many parasites and bacteria live in fresh water and can cause serious illness. If you are unsure about the safety in the area you are visiting, check with a reliable source before swimming. Properly chlorinated pools and salt water are generally safe from infectious diseases.

OTHER HEALTH CONCERNS

COVID 19:

COVID-19 is a respiratory virus that is spread through direct contact with an infected person as well as through respiratory droplets produced when an infected person coughs or sneezes. Symptoms of COVID-19 may appear 2-14 days after exposure and may include fever, cough, and shortness of breath. The illness can also cause muscle or body aches, sore throat, vomiting and diarrhea. Reported illnesses have ranged from mild symptoms to severe illness and death.

It's recommended all eligible travelers should be up to date with their COVID-19 vaccines which is the best way to prevent infection. Please see Your COVID-19 Vaccination for more information.

SIT recommends that students check the country's consular website for up-to-date information on entry and exit requirements.

Other precautions to minimize your exposure to Covid include:

- Clean your hands frequently with soap and water or alcohol-based disinfectant.
- Wear an N-95 mask when in large crowds or on public transportation.
- Avoid people with suspected or confirmed Covid infection.
- Inform your Program staff immediately if you have a respiratory illness; have a fever or are feeling sick; if you have been in close contact with a person known to have COVID-19; and/or have recently traveled from an area with widespread or ongoing community spread of COVID-19.
- Avoid travel if you are sick or have a fever. Your Program staff will make appropriate accommodations for students who are ill.

- Wear an N-95 facemask, especially when you are around other people (e.g., sharing a room or vehicle) and before you enter a health care provider's office.

Blood Borne Pathogens:

Pathogens such as Hepatitis B and C and HIV are more prevalent in certain parts of the world and can cause severe, long-term illness. Practices to minimize transmission through sanitization may be less regulated than at home.

You can minimize your risk by:

- Practicing safe sex, including the use of condoms
- Not sharing needles, syringes, razors, toothbrushes, or medical equipment, such as a glucose monitor.
- Not getting tattoos while traveling overseas.

Tuberculosis:

Tuberculosis (TB) is a bacterial disease spread by airborne droplets from an infected person. Bacteria in the lungs can move through the blood to infect other parts of the body, such as the kidney, spine, and brain. TB occurs throughout the world but is much more common in some countries including CHILE. Travelers can protect themselves by avoiding close contact with people who are coughing and who look sick. If you believe you were exposed, you should ask for TB testing were you return home.

Altitude:

You may be visiting locations at high altitudes. Even healthy, athletic individuals may become ill at altitudes over 10,000 ft. Common symptoms are unbearable headache and severe shortness of breath out of proportion to the mild fatigue most people experience while becoming acclimatized. Ascending gradually and resting during the first few days can minimize the risk of altitude sickness.

Individuals with lung disorders, such as asthma, heart concerns and any other preexisting medical condition including sickle cell or diabetes should consult a health care provider before traveling to high altitudes. If your health care provider

WORLD LEARNING PROGRAM HEALTH GUIDELINES & REQUIREMENTS: CHILE

has given approval for high altitude travel, please reach out to the Student Health Office so that we can advise the Program staff and our local health care providers accordingly. Bring full medical notes with you to assist local health care providers in case of need.

You should consult your health care provider about taking acetazolamide (Diamox). Please note that alcohol and sedatives have greater effect at high altitudes. Any symptoms of severe altitude illness should result in immediate descent.

Sun Exposure:

World Learning recommends the use of sunglasses, wide-brimmed hats, sunscreen lotions, and lip protection to reduce problems related to sun exposure.

Air Quality:

Students with a history of asthma, other lung disease or allergies should be warned that air pollution may be worse than you are used to, causing more symptoms while you are traveling. Asthmatics should carry emergency medicines for severe asthma attacks.

Dehydration:

Dehydration occurs when the body is losing fluid faster than it can be replaced. This can be extreme in cases of diarrhea or vomiting, or gradual in cases of exertion without adequate fluid intake especially in hot climates and at high altitude. Travelers should drink safe water or other safe beverages regularly and should avoid excessive sugary or caffeinated beverages as these can increase fluid loss. Consistent fluid consumption throughout the day is best.

World Learning is committed to supporting the emotional and physical wellbeing of all students. If you have any questions or concerns about any of these health guidelines, please reach out to the Student Health Office associated with your upcoming Program.

